CORRECTION

Respiratory Research



Correction: Combination of betulinic acid and EGFR-TKIs exerts synergistic anti-tumor effects against wild-type EGFR NSCLC by inducing autophagy-related cell death via EGFR signaling

Han Wang^{1,2†}, Xiaohui Du^{1†}, Wenwen Liu¹, Congcong Zhang¹, Ying Li¹, Jingwen Hou¹, Yi Yu¹, Guiru Li^{1*} and Qi Wang^{1*}

Correction to: Respiratory Research (2024) 25:215 https://doi.org/10.1186/s12931-024-02844-9

In the original publication of this article [1], there was an error in Fig. 5. The western blots lines reporting the EGFR and p-Her2 bands for H1299 cell lines were duplicated. The EGFR bands was mistakenly repeated in p-Her2 bands. For completeness and transparency, the old incorrect and correct Fig. 5 are displayed in this correction article.

[†]Han Wang and Xiaohui Du contributed equally to this study.

The original article can be found online at https://doi.org/10.1186/s1 2931-024-02844-9.

*Correspondence: Guiru Li Liguiru_dlmu@126.com Qi Wang wqdlmu@163.com ¹The Second Hospital of Dalian Medical University, Dalian 116023, China ²Guangzhou women and children's medical center, Guangzhou Medical University, Guangzhou 510623, China



© The Author(s) 2025. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/. The Creative Commons Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Incorrect Fig. 5:



Fig. 5 Combination treatment enhanced the suppression of EGFR and its downstream signaling and bypass pathways in wt-EGFR NSCLC cells. A549 and H1299 cells were treated with DMSO, BA, gefitinib, osimertinib or the indicated combination and then harvested for preparation of whole-cell protein lysates and subsequent western blotting to detect the indicated proteins

Correct Fig. 5:



Fig. 5 Combination treatment enhanced the suppression of EGFR and its downstream signaling and bypass pathways in wt-EGFR NSCLC cells. A549 and H1299 cells were treated with DMSO, BA, gefitinib, osimertinib or the indicated combination and then harvested for preparation of whole-cell protein lysates and subsequent western blotting to detect the indicated proteins

Published online: 13 April 2025

References

 Wang H, Du X, Liu W, et al. Combination of betulinic acid and EGFR-TKIs exerts synergistic anti-tumor effects against wild-type EGFR NSCLC by inducing autophagy-related cell death via EGFR signaling pathway. Respir Res. 2024;25:215.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.